

# Dosing & monitoring

**Intended for patients prescribed Duvyzat® ▼ (givinostat)**  
Including parents, guardians and caregivers

This medicine is subject to additional monitoring. Information about reporting side effects can be found on the back page.  
If you have any questions about dosing or administering Duvyzat®, please speak with your healthcare team.

All photographic images  
within this booklet  
feature real patients.  
They are shown for  
illustrative purposes  
only, and individual  
experiences may vary



**Make a note of your  
Duvyzat® dose and keep  
track of appointments**



# Your recommended dose



**Your doctor will decide the correct dose based on body weight and other factors. Do not adjust the dose yourself.**

Always take this medicine exactly as your doctor, pharmacist or nurse has told you. Check with your doctor, pharmacist or nurse if you are not sure. The recommended dose of Duvyzat<sup>®</sup> depends on your body weight.

## Your body weight:

Blank space for your doctor to write in body weight

## Your Duvyzat<sup>®</sup> recommended dose:

## Your Duvyzat<sup>®</sup> modification dose 1:

## Your Duvyzat<sup>®</sup> modification dose 2:

A detailed method of administration can be found in the 'Dosing instructions' leaflet, available as a PDF at: [myeveryday.support](https://myeveryday.support)



### If you take more Duvyzat<sup>®</sup> than you should.

Contact your doctor or a hospital if you take more than the prescribed dose of Duvyzat<sup>®</sup>.



### If you forget a dose, take the next dose when planned.

Do not take a double dose.



### Do not stop taking Duvyzat<sup>®</sup> without talking to your doctor.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

**Please take this card to all your appointments so the results can be added, and future appointment dates can be checked and confirmed.**

# Your monitoring calendar



**Regular monitoring is essential when treated with Duvizat®. Below is a table for you to keep track of appointments**

	Platelets		Triglycerides		Body weight			
<b>At the start of treatment</b>	Date:		Date:		Date:			
	Result:		Result:		Result:			
<b>Month 1: Week 2</b>	Date:		Date:		Date:			
	Result:		Result:		Result:			
<b>Month 1: Week 4</b>	Date:		Month 1 triglycerides test can take place any time during month 1		Date:			
	Result:				Result:			
<b>Month 2: Week 6</b>	Date:				Date:			
	Result:				Result:			
<b>Month 2: Week 8</b>	Date:				Date:			
	Result:				Result:			
<b>Month 3</b>	Date:				Date:		Date:	
	Result:				Result:		Result:	
<b>Month 6</b>	Date:		Date:		Date:			
	Result:		Result:		Result:			
<b>Ongoing</b>	Every 3 months		Every 6 months		As required			

## It is important to monitor your platelets and triglycerides to help prevent complications

Duvizat® can cause dose-related thrombocytopenia and other signs of myelosuppression, including decreased haemoglobin and neutropenia. Duvizat® can also cause elevations in triglycerides.

### Additional monitoring

Your healthcare team may recommend additional ECG monitoring due to underlying cardiac disease or other medications you are taking.

Further information can be found in the 'Potential side effects' leaflet, available as a PDF at: [myeveryday.support](https://myeveryday.support)

Date of preparation: March 2026

Job Code: GB-DVZ-26-00005



## Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in the package leaflet. You can also report side effects directly via the Yellow Card Scheme at: <https://yellowcard.mhra.gov.uk>

By reporting side effects, you can help provide more information on the safety of this medicine. This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See <https://yellowcard.mhra.gov.uk> for how to report side effects.