

Potential side effects

Intended for patients prescribed Duvyzat® ▼ (givinostat)
Including parents, guardians and caregivers

This medicine is subject to additional monitoring. Information about reporting side effects can be found on the back page.
If you have any questions about dosing or administering Duvyzat®, please speak with your healthcare team.

All photographic images within this booklet feature real patients. They are shown for illustrative purposes only, and individual experiences may vary

Make yourself aware of the known and potential side effects of Duvyzat®




Known side effects of Duvyzat®



Unfortunately, there's no way to predict whether side effects will occur

Both you and family members should watch out for potential signs, and inform your healthcare team if any issues arise.

Known side effects:



Reduced platelet count (thrombocytopenia)

If you have unusual bleeding or small red to purple spots on the skin, inform your healthcare team immediately.



Diarrhoea

If you experience diarrhoea, keep track of how often it occurs and how severe it is, and contact your healthcare team. It is important to stay well hydrated.



Increased blood fat levels (elevated triglycerides)

These changes in the blood initially show no symptoms but are detected early thanks to regular monitoring.



Changes in heart rhythm (QTc prolongation)

Some people may experience changes in the heart's rhythm. This usually causes no symptoms, but your healthcare team may monitor this with heart tests called ECGs.

Important note

Serious side effects can also occur during treatment with Duvyzat®. On this basis, your healthcare team will carry out regular tests and take appropriate action if any abnormalities are detected.



Potential side effects



Additional side effects may occur whilst taking Duvyzat® - some are very common and some common

Very common (may affect more than 1 in 10 people)

- abdominal pain
- decrease in blood platelet count (thrombocytopenia)
- diarrhoea
- elevated levels of blood fats (triglycerides)
- fever
- nausea
- vomiting

Common (may affect up to 1 in 10 people)

- constipation
- decreased appetite
- fatigue
- joint pain (arthralgia)
- muscle pain (myalgia)
- rash

Not known: frequency cannot be estimated from the available data:
hypothyroidism and/or increased in thyroid stimulating hormone

Important note

Serious side effects can also occur during treatment with Duvyzat®. On this basis, your healthcare team will carry out regular tests and take appropriate action if any abnormalities are detected.



Date of preparation: March 2026

Job Code: GB-DVZ-26-00006



Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in the package leaflet. You can also report side effects directly via the Yellow Card Scheme at: <https://yellowcard.mhra.gov.uk>

By reporting side effects, you can help provide more information on the safety of this medicine. This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See <https://yellowcard.mhra.gov.uk> for how to report side effects.